

Effects of elevated carbon dioxide on arbuscular mycorrhizal fungi activities and soil microbial properties in soybean (*Glycine max* L. Merrill) rhizosphere

Nurudeen Olatunbosun Adeyemi^{1*}, Muftau Olaoye Atayese¹, Michael Dare², Adebanke Olubode²

¹Federal University of Agriculture, Abeokuta, College of Plant Science and Crop Production, Department of Plant Physiology and Crop Production, Alabata, Ogun State, Nigeria

²Federal University of Agriculture, Abeokuta, College of Plant Science and Crop Production, Department of Soil Science and Land Management, Alabata, Ogun State, Nigeria

Article Details: Received: 2020-03-31 | Accepted: 2020-04-28 | Available online: 2020-09-30

<https://doi.org/10.15414/afz.2020.23.03.109-116>



Licensed under a Creative Commons Attribution 4.0 International License



Arbuscular mycorrhizal fungi (AMF) help in promoting plant growth and mediating key belowground processes, however, AMF responses to the continuous increase in the atmospheric carbon dioxide (CO₂) is yet elusive. This has led to considerable interest in the impacts elevated CO₂ on AMF and belowground processes in recent years. The present study investigated the effect of elevated CO₂ on AMF sporulation and root colonization and soil microbial properties in the rhizosphere of soybean. The pot experiment consisted of two levels of CO₂ (ambient; 350 ppm and elevated; 550 ppm) and three soybean cultivars (TGx 1440-1E, TGx 1448-2F and TGx 1480-2F) conducted in open top chambers, laid out in randomized complete block design, replicated thrice. The results showed that elevated CO₂ increased the AMF spore density and root colonization of the soybean cultivars. Elevated CO₂ increased the microbial biomass carbon (34.2–45.4%), microbial biomass nitrogen (44.6–54.9%), soil nitrogen (30.3–50.6%), available phosphorus (20.8–45.7%) in the rhizosphere of the soybean cultivars compared to the ambient CO₂. These could have resulted in increased plant biomass, pod number, 100-seed weight and seed yield under elevated CO₂. From the results of this study, increased atmospheric CO₂ regulates AMF activities, microbial properties and improve soybean performance. Thus, this study may help to a better understanding of the responses of AMF and belowground process with increasing atmospheric CO₂.

Keywords: arbuscular mycorrhizal fungi, climate change, CO₂ enrichment, microbial biomass, open top chambers

1 Introduction

The atmospheric carbon dioxide (CO₂) concentration is constantly increasing since industrial revolution due to anthropogenic activities, and is expected to double by 2050 (IPCC, 2013; Cotton et al., 2015; Kumar et al., 2019). Indeed, elevated CO₂ directly increases the net primary photosynthesis of crops, particularly C₃ plants; stimulating the production of aboveground biomass production (Oliveira et al., 2010) and carbon (C) fluxes from aboveground parts into soil (Ghannoum et al., 2010; McCarthy et al., 2010). Thus, the continuous rising atmospheric CO₂ concentrations may indirectly affect the responses of soil microbial communities and the associated C and nitrogen (N) cycling in the rhizosphere (Bardgett et al., 2008; Butterly et al., 2016). Since plant

mediates belowground processes, the responses of associated root microbes, which serve as sinks for the photoassimilates are likely to be system specific (Carrillo et al., 2014; Goicoechea et al., 2014). Given that soil microbes, particularly arbuscular mycorrhizal fungi (AMF) are actively involved in C and nutrient cycling in the agroecosystems (Cairney, 2012), there is an urgent need to better understand the effects of elevated CO₂ on AMF and soil microbial properties in the rhizosphere of agricultural crops.

AMF are one of the most important soil microbes forming symbiotic association with 90% of terrestrial plants including most agricultural crops (Brundrett, 2009). AMF play an important role in promoting plant growth through nutrient uptake, especially phosphorus, in exchange for

*Corresponding Author: Nurudeen Olatunbosun Adeyemi, Federal University of Agriculture, Abeokuta, College of plant Science and Crop production, Department of Plant Physiology and Crop Production, P.M.B.2240, Alabata, Ogun State, Nigeria. E-mail: adeyemisworld@gmail.com. ORCID: 0000-0001-6341-775X

carbon from the host plant (Wipf et al., 2019). They also account for about 5–50% of microbial biomass and about 80% of fungal biomass (Kabir et al., 1997). The impacts of elevated CO₂ on AMF are often reported by increased root colonisation, which is usually attributed to increased C assimilation rates by the host plant in exchange for increased nutrients demand by the host plant (Johnson and Gehring, 2007; Cairney, 2012). However, the shifts towards greater root colonisation by AMF may have important consequences on soil properties including nutrient supply to the plant, particularly P and N and microbial biomass C and N (Drigo et al., 2013).

Soybean [*Glycine max* (L.) Merrill] like most other legumes is highly mycorrhizal dependent (Adeyemi et al., 2020). Being a C₃ plant and a crop with high phosphorus (P) demand may be more sensitive to elevated CO₂ compared to other non-leguminous crops (Rogers et al., 2006). Soybean is reported to respond positively to elevated CO₂ with higher productivity (Ziska and Bunce, 2000; Sakariyawo et al., 2016; Adeyemi et al., 2017). Previous researchers have focused their studies on above ground changes in response to elevated CO₂, but recently some of the findings revealed that elevated CO₂ could also influence below ground microbial properties, which in turn affects plant productivity (Bhattacharyya et al., 2016). In addition, the alteration of AMF responses in terms of sporulation and colonization of plant roots under elevated CO₂ are yet elusive and there are inconsistencies in previous reports (Johnson et al., 2005; Cotton et al., 2015; Parameswaran et al., 2019). Thus, this could hinder the prediction of microbial responses, particularly AMF in agroecosystem functioning (Singh et al., 2011; Johnson et al., 2013). Therefore, the growing awareness of the significant role of AMF for the maintenance of agroecosystem and promoting plant growth has led to considerable interest in the impacts of elevated CO₂ on AMF activities in this present study.

Thus, this study aims to investigate the effects of short-term elevated CO₂ on AMF sporulation and root colonization, soil microbial properties in roots of soybean cultivars grown in open top chambers. The results of this study may help to gain insights into how AMF and belowground process respond to the increasing atmospheric CO₂.

2 Materials and methods

2.1 Experimental treatments and design

The study was conducted in Open Top Chambers (OTCs) at College of Plant Science and Crop Production, Federal University of Agriculture, Abeokuta. The pot experiment consisted of three promiscuous soybean cultivars (TGx 1448-2E, TGx 1440-1E and TGx 1740- 2F) grown at two CO₂

levels (ambient; 350 ppm and elevated; 550 ppm) laid out in completely randomised design, replicated thrice. The soil used for the experiment was a non-sterile obtained from the Teaching and Research Farms of the University, with textural soil of sandy (Table 1). Each pot was filled with 10 kg of soil. The soil in the pots was watered to 60% of the water-holding capacity. Soybean cultivars TGx 1448-2E, TGx 1440-1E and TGx 1740- 2F) were obtained from International Institute of Tropical Agriculture (IITA), Ibadan. Three seeds were sown in each pot on 10th of July, 2015 and emerged seedlings were thinned to two plants per pot. As plants grew bigger, pots were watered to 70% of water-holding capacity. The pots were placed in OTCs at 350 and 550 ppm CO₂ Carbon dioxide enrichment

Table 1 Soil physical and chemical properties of the experimental site

Soil Property	Value
Texture	Sand
Sand (%)	87.9
Silt (%)	7.49
Clay (%)	4.61
pH (H ₂ O)	6.5
Organic matter (%)	1.12
Nitrogen (mg kg ⁻¹)	0.80
Available phosphorus (mg kg ⁻¹)	5.23
Potassium (cmol kg ⁻¹)	0.44
Calcium (cmol kg ⁻¹)	2.54
Magnesium (cmol kg ⁻¹)	0.73
Sodium (cmol kg ⁻¹)	0.29
Total exchangeable acidity (cmol kg ⁻¹)	0.11
Exchangeable cation exchange capacity (cmol kg ⁻¹)	3.92

Each OTC was 3 × 4 m in cross section and 2.5 m in height. The frame was made of pipes covered with transparent polyvinyl chloride nylon sheet (transparency 85%). CO₂ was produced using a modified method of Saitoh et al. (2004). Yeasts and sugar were used to generate CO₂. Twelve CO₂ generator bottles were placed in the OTC to elevate the CO₂ concentration. The maximum and minimum CO₂ concentration in the OTC was measured daily throughout the enrichment period using a portable CO₂ meter [NDIR Gas Analyzer (Bentech GM8883), China]. To avoid chamber specific bias in the experiment, pots were rearranged in the OTCs every week.

2.2 Data collection

Estimation of AMF spore density and root length colonisation

Rhizosphere soil samples from 0–20 cm depth were collected at 20, 40 and 60 days after planting (DAP) to determine AMF spore density. Extraction of AMF spores from the soil sample was conducted using the modified wet sieving method of Giovannetti and Mosse (1980), followed by centrifugation in 40% sucrose solution to extract the spores. Extracted spores were counted under a dissecting microscope at $\times 100$ magnification.

Collected root samples were prepared using the method as described by Phillips and Hayman (1970). Root samples collected were bleached in 10% KOH at 90 °C for 30 minutes and stained in acidic glycerol containing trypan blue lacto-glycerol (1 : 1 : 1 : 0.5 g) at 90 °C for 2 minutes. The roots were viewed under compound microscope to determine percent root colonization using the formula below as described by Adeyemi et al. (2019, 2020):

$$\text{percent root colonisation} = \frac{\text{number of root infected}}{\text{total number of roots}} \times 100$$

Measurement of soil microbial biomass C and N

Soil microbial biomass C and N were measured by using the chloroform fumigation extraction method (Vance et al., 1987). The dry soil samples were divided into two portions (10 g per portion) to measure the soil microbial biomass carbon (SMBC) and soil microbial biomass nitrogen (SMBN). SMBC in filtrates was determined by the $K_2Cr_2O_7$ wet-oxidation method. Briefly, 2 ml of extract, 2 ml of 1 N $K_2Cr_2O_7$ and 4 ml of 98% H_2SO_4 were reacted at 130 °C for 40 min. Solutions with known concentrations of $C_{12}H_{22}O_{11}$ were included as standards. After cooling, the C concentration in digests was determined spectrophotometrically at 600 nm. The C contained within digested non-fumigated samples

was denoted extractable organic C and the SMBC was estimated as the difference between the fumigated and non-fumigated samples and applying a conversion factor of 0.38 (Singh et al., 2017). SMBN in filtrates was assayed by the semi-micro Kjeldahl method. Specifically, 2.5 ml of extract and digestion mix (50 g $K_2S_2O_8$ and 30 g H_3BO_4 in 100 ml of 3.75 M NaOH) were autoclaved (121 °C, 104 kPa) for 30 min and stored at 4 °C until analysis. Solutions with known concentrations of $CO(NH_2)_2$ were included as controls. SMBN was calculated from the differences between the organic N extracted from the fumigated and non-fumigated samples by using conversion factor of 0.54 (Huang et al., 2014).

2.3 Soybean growth

The plants were harvested and oven dried (60 °C, 72 h) prior to weighing to determine dry plant biomass. At 95% harvest maturity, yield attributes and seed yield were collected on the number of pods per plant, number of seeds per pod, 100 seed weight and seed yield per plant.

2.4 Statistical analysis

Data were subjected to analysis of variance (ANOVA). The Duncan's multiple range (DMR) test at $P < 0.05$ probability level was used to determine the differences among mean values. The statistical package used was Genstat 12th Edition.

3 Results and discussion

3.1 Effects of elevated CO_2 on AMF spore density and percent root colonization

The result showed that elevated CO_2 significantly increased the spore density of AMF in rhizospheric soil of the soybean cultivars at 40 and 60 DAP (Table 2). Higher spore density of 166 and 216 spores per 20 g soil was recorded in TGx 1448 2E grown under elevated CO_2 concentration (500 ppm) at 40 and 60 DAS respectively. The percent root colonization of the soybean cultivars

Table 2 Effect of elevated CO_2 on AMF spore density in rhizospheric soil of three soybean cultivars

Cultivars	Atmospheric CO_2	AMF spore density (per 20 g dry soil)		
		20 DAP	40 DAP	60 DAP
TGx 1440 1E	Ambient	30.0 ^a	111.7 ^{cd}	143.3 ^b
	Elevated	38.3 ^a	155.0 ^{ab}	208.3 ^a
TGx 1448 2E	Ambient	33.3 ^a	115.0 ^c	155.0 ^b
	Elevated	35.0 ^a	166.7 ^a	216.7 ^a
TGx 1740 2F	Ambient	26.7 ^a	86.7 ^d	135.0 ^b
	Elevated	36.3 ^a	128.3 ^{bc}	211.7 ^a

Treatments followed by the same letters are not significantly different at $P < 0.05$ level; DAP (days after planting)

was significantly influenced by elevated CO₂ with higher root colonization observed in the cultivars grown under elevated CO₂ (Figure 1a, b, c). The mean percent colonization ranged from 50.7 to 88% at 60 DAP.




Figure 1 Effect of elevated CO₂ on AMF root colonization of soybean cultivars at 20 DAP (A), 40 DAP (B) and 60 DAP (C)

3.2 Soil microbial biomass C and N

Elevated CO₂ caused significant changes in the soil microbial biomass C and N. Specifically, the SMBC increased significantly under the elevated CO₂ in the rhizospheric soils of the three soybean cultivars. The SMBC in the soil layer of the soybean was in the range of 511 to 659 mg kg⁻¹ under the elevated CO₂ compared to range of 336 to 360 mg kg⁻¹ under ambient condition. Compared with the ambient treatment, SMBC significantly increased by 40.9%, 45.4%, and 34.2% in the roots of TGx 1440 1E, TGx 1448 2E and TGx 1740 2F cultivars respectively (Figure 2). Moreover, SMBC was higher in TGx 1448 2E than other cultivars under the elevated CO₂. Similarly, SMBN increased with an increased CO₂ (Figure 3). Compared with ambient CO₂,

SMBN significantly increased by 44.6%, 50.8% and 54.9% in the roots of TGx 1440 1E, TGx 1448 2E and TGx 1740 2F cultivars respectively.




Figure 2 Dynamics of soil microbial biomass carbon (SMBC) in the 0–10 cm soil layer of soybean cultivars as affected by elevated CO₂




Figure 3 Dynamics of soil microbial biomass nitrogen (SMBN) in the 0–10 cm soil layer of soybean cultivars as affected by elevated CO₂

3.3 Soil nitrogen and available phosphorus

The result showed that the total nitrogen and available phosphorus in the soil increased significantly with elevated CO₂ in the soil of all the three cultivars. Compared with ambient CO₂, total N significantly increased by




Figure 4 Dynamics of soil total nitrogen (g kg⁻¹) in the 0–10 cm soil layer of soybean cultivars as affected by elevated CO₂

37.2%, 50.6% and 30.3% in the soil of TGx 1440 1E, TGx 1448 2E and TGx 1740 2F cultivars respectively (Figure 4). Similarly, compared with ambient CO₂, the soil available P significantly increased by 45.7%, 20.8% and 39.8% in the roots of TGx 1440 1E, TGx 1448 2E and TGx 1740 2F cultivars respectively (Figure 5).




Figure 5 Dynamics of soil available phosphorus (mg kg⁻¹) in the 0–10 cm soil layer of soybean cultivars as affected by elevated CO₂

3.4 Effects of elevated CO₂ on plant biomass and yield components of soybean cultivars

The result showed that the above ground dry biomass and all the yield attributes measured were significantly influenced by elevated CO₂ (Table 3). The soybean dry biomass, number of pod, number of seed per pod, 100-seed weight and seed yield per plant were significantly higher in the three cultivars grown under elevated CO₂ compared to the ambient control.

The effects of the elevated CO₂ on soil microorganisms may be direct or indirect. This present study investigated the effects of elevated CO₂ on AMF sporulation and root colonization and soil microbial properties in roots of soybean cultivars. In the present study, root colonization of the soybean cultivars increased under elevated CO₂. Elevated CO₂ concentrations have been reported to

affect the AMF hyphal growth and root colonization. The result of this study agrees with reports of previous studies (Compant et al., 2010). However, Goicoechea et al. (2014), have also reported a contrary result. The increased root colonization of the soybean cultivars may be explained by increase flow of C from the host to AMF being a strong sink of C fixed during photosynthesis by host plants under elevated CO₂ (Smith and Read, 2008; Ghannoum et al., 2010; Cairney, 2012) in exchange for additional nutrients uptake, especially P (Moran and Jastrow, 2010). For example, Rillig and Allen (1999) reported an increased growth of external as well as internal hyphae of AMF in the rhizosphere of *Prunella vulgaris* under elevated CO₂ (600 ppm) due to increased root biomass and higher allocation of fixed C to the external hyphae. Furthermore, in a meta-analysis by Alberton et al. (2005), a significant positive response of AMF to elevated CO₂, causing a 21% increase in root colonization was revealed. However, Gavito et al. (2000) observed no effect of elevated CO₂ (700 ppm) on root colonization in pea (*Pisum sativum* L.). Another mechanism for the increased root colonization of the soybean cultivars could be attributed to rapid release of flavonols needed for chemotactic action on AMF, which stimulate the hyphal growth and root colonization (Bécard et al., 1992).

In this study, microbial biomass C and N increased under elevated CO₂, which is in agreement with some elevated CO₂ studies (Jin et al., 2013; Butterly et al., 2016; Panneerselvam et al., 2019). The increase in the microbial biomass C have been associated with higher C inputs or rhizodeposition (Drissner et al., 2007; Cheng et al., 2011), which stimulates microbial growth and increases biomass. A recent review of 68 studies revealed that the average microbial biomass C increase is 14% and N is 7.4% (Liu et al., 2018). Thus, the effects depend on the specific soil conditions, which are mainly controlled by N limitation (Sillen and Dieleman, 2012). However, elevated CO₂ did not influence the microbial biomass C in some other studies (Weigel et al., 2005; Reinsch et al., 2013;

Table 3 Effect of elevated CO₂ on dry biomass, pod number, seed number per pod, seed weight 100 grain⁻¹, and seed weight per plant of soybean cultivars

Cultivars	Atmospheric CO ₂	Dry biomass (g plant ⁻¹)	Pod number (n plant ⁻¹)	Seed number (n pod ⁻¹)	Seed weight (g 100 grain ⁻¹)	Seed yield (g plant ⁻¹)
TGx 1440 1E	Ambient	24.7 ^d	41.0 ^c	2.4 ^c	9.73 ^b	9.69 ^d
	Elevated	38.4 ^{ab}	57.0 ^a	2.7 ^{ab}	11.3 ^a	17.5 ^b
TGx 1448 2E	Ambient	30.9 ^c	48.4 ^b	2.6 ^b	9.9 ^b	12.4 ^c
	Elevated	41.3 ^a	61.0 ^a	2.8 ^a	11.4 ^a	19.5 ^a
TGx 1740 2F	Ambient	19.7 ^d	31.6 ^d	2.3 ^c	8.22 ^b	6.13 ^e
	Elevated	35.2 ^b	55.2 ^a	2.7 ^{ab}	10.6 ^a	15.8 ^b

Treatments followed by the same letters are not significantly different at $P < 0.05$ level

Haugwitz et al., 2014). The increased microbial biomass N observed in this study could be associated with the activities of the rhizobia in the roots of the soybean, which helps in N-fixation in the rhizosphere (Kuzyakov et al., 2018). However, in some studies, microbial biomass N was not affected by elevated CO₂ (Schortemeyer et al., 2000; Zak et al., 2000). Furthermore, the higher soil N in this study may be attributed to reduced nitrate leaching under elevated CO₂ (Cannell and Thornley, 1998; Matamala and Drake, 1999). In contrary, elevated CO₂ has been observed to increase denitrification or N₂O fluxes, therefore, potentially reduce N retention (Ineson et al., 1995). Generally, the increased microbial biomass properties (C and N), soil N and P could also be explained by the functional role of AMF for nutrients cycling particularly C, N and P, which is more intensified under elevated CO₂ (Denef et al., 2007; Drigo et al., 2013; Fang et al., 2015). Thus, the microbial responses to elevated CO₂ are expected to vary among ecosystems, since responses to elevated CO₂ are positively related to the plants or crops grown, soil-type specific and microbial communities in the soil (Nie et al., 2013; Pendall et al., 2013; Procter et al., 2014),

The present study also revealed increased seed yield and yield attributes of the soybean cultivars under elevated CO₂. The result was in consistent with previous reports (Sakariyawo et al., 2016; Adeyemi et al., 2017). This positive response could be attributed to enhance net photosynthesis and dry matter accumulation prior to the formation of the yield attributes, resulting in improve seed yield. Results of several other studies have suggested that elevated CO₂ will result in increased photosynthesis of C₃ plants by 30–50% (Yang et al., 2006; Ainsworth and Rogers, 2007). In addition, the increased yield performance in this study could also be attributed to the enhanced AMF root colonization, which helps in nutrients uptake, particularly P and water, as well as improved soil properties in terms of soil N and P and microbial C and N.

4 Conclusions

The results of this study demonstrates that elevated CO₂ concentration had significant effect on AMF activities in the rhizosphere as well as the soil microbial biomass by increasing the root colonization, soil N and P and microbial biomass C and N. This resulted in higher seed yield and yield attributes of the soybean cultivars. Further investigation is required to ascertain the responses of AMF and other soil microbial communities in the cycling of C, N and P under different crops in response to elevated CO₂. This study may help to better forecast how the AMF and soil microbes respond with increasing atmospheric CO₂.

References

- ADEYEMI, N.O. et al. (2020). Effect of commercial arbuscular mycorrhizal fungi inoculant on growth and yield of soybean under controlled and natural field conditions. *Journal of Plant Nutrition*, 43(4), 487–499. <https://doi.org/10.1080/01904167.2019.1685101>
- ADEYEMI, N.O. et al. (2019). Identification and relative abundance of native arbuscular mycorrhizal fungi associated with oil-seed crops and maize (*Zea mays* L.) in derived savannah of Nigeria. *Acta fytotechn zootechn*, 22(3), 84–89. <https://doi.org/10.15414/afz.2019.22.03.84-89>
- ADEYEMI, N., SAKARIYAWO, O. and ATAYESE, M. (2017). Yield and yield attributes responses of soybean (*Glycine max* L. Merrill) to elevated CO₂ and arbuscular mycorrhizal fungi inoculation in the humid transitory rainforest. *Notulae Scientiae Biologicae*, 9(2), 233–241. <https://doi.org/10.15835/nsb9210002>
- AINSWORTH, E. A. and ROGERS, A. (2007). The response of photosynthesis and stomatal conductance to rising [CO₂]: Mechanisms and environmental interactions. *Plant, Cell and Environment*, 30, 258–270. <https://doi.org/10.1111/j.1365-3040.2007.01641.x>
- ALBERTON, O., KUYPER, T.W. and GORISSEN, A. (2005). Taking mycocentrism seriously: mycorrhizal fungal and plant responses to elevated CO₂. *NewPhytol.*, 167, 859–868
- BARDGETT, R.D., FREEMAN, C. and OSTLE, N.J. (2008). Microbial contributions to climate change through carbon cycle feedbacks. *ISME J.*, 2, 805–814.
- BÉCARD, G. et al. (1992). Extensive *in vitro* hyphal growth of vesicular-arbuscular mycorrhizal fungi in presence of CO₂ and flavenols. *Applied Environmental Microbiology*, 58, 821–825.
- BHATTACHARYYA, P. et al. (2016). Elucidation of rice rhizosphere metagenome in relation to methane and nitrogen metabolism under elevated carbon dioxide and temperature using whole genome metagenomic approach. *Sci. Total Environ*, 542, 886–898.
- BRUNDRETT, M.C. (2009). Mycorrhizal associations and other means of nutrition of vascular plants: understanding the global diversity of host plants by resolving conflicting information and developing reliable means of diagnosis. *Plant Soil*, 320, 37–77.
- BUTTERLY C.R. et al. (2016). Long-term effects of elevated CO₂ on carbon and nitrogen functional capacity of microbial communities in three contrasting soils. *Soil Biology and Biochemistry*, 97, 157–167. <https://doi.org/10.1016/j.soilbio.2016.03.010>
- CAIRNEY, J.W.G. (2012). Extramatrical mycelia of ectomycorrhizal fungi as moderators of carbon dynamics in forest soil. *Soil Biology and Biochemistry*, 95, 198–208.
- CANNELL, M.G.R. and THORNLEY, H.M. (1998). N-poor ecosystems may respond more to elevated [CO₂] than N-rich ones in the long term. A model analysis of grassland. *Glob. Change Biol.*, 4, 431–442.
- CARRILLO, Y. et al. (2014). Plant rhizosphere influence on microbial C metabolism: the role of elevated CO₂, N availability and root stoichiometry. *Biogeochemistry*, 117, 229–240. <https://doi.org/10.1007/s10533-014-9954-5>
- CHENG, L. et al. (2011). Soil microbial responses to elevated CO₂ and O₃ in a nitrogen-aggrading agroecosystem. *Plos One*, 6, e21377. <https://doi.org/10.1371/journal.pone.0021377>

- COMPANT, S. et al. (2010). Climate change effects on beneficial plant–microorganism interactions. *Microbiology Ecology*, 73, 197–214.
- COTTON, T.E. et al. (2015). Fungi in the future: interannual variation and effects of atmospheric change on arbuscular mycorrhizal fungal communities. *New Phytol.*, 205, (4), 1598–1607. <https://dx.doi.org/10.1111/nph.13224>
- DENEFF, K. et al. (2007). Community shifts and carbon translocation within metabolically-active rhizosphere microorganisms in grasslands under elevated CO₂. *Biogeosciences*, 4, 769–779
- DRIGO, B. et al. (2013). Impacts of 3 years of elevated atmospheric CO₂ on rhizosphere carbon flow and microbial community dynamics. *Glob. Change Biol.*, 19(2), 621–636. <https://doi.org/10.1111/gcb.12045>
- DRISSNER, D. et al. (2007). Nine years of enriched CO₂ changes the function and structural diversity of soil microorganisms in a Grassland. *Eur. J. Soil Sci.*, 58, 260–269.
- FANG, H.J. et al. (2015). Elevated atmospheric carbon dioxide concentration stimulates soil microbial activity and impacts water-extractable organic carbon in an agricultural soil. *Biogeochemistry*, 122, 253–267. <https://doi.org/10.1007/s10533-014-0039-2>
- GAVITO M.E. et al. (2000) Atmospheric CO₂ and mycorrhiza effects on biomass allocation and nutrient uptake of nodulated pea (*Pisum sativum* L.) plants. *J Exp Bot*, 51, 1931–1938.
- GHANNOUM, O. et al. (2010). Exposure to preindustrial, current and future atmospheric CO₂ and temperature differentially affects growth and photosynthesis in *Eucalyptus*. *Global Change Biology*, 16, 303–319.
- GIOVANETTI, M. and MOSSE, B. (1980). An evaluation of techniques for measuring vesicular arbuscular mycorrhizal infection in roots. *New Phytologist*, 84, 489–500.
- GOICOECHEA, N. et al. (2014) Increased photosynthetic acclimation in alfalfa associated with arbuscular mycorrhizal fungi (AMF) and cultivated in greenhouse under elevated CO₂. *Journal of Plant Physiology*, 171(18), 1774–1781. <https://doi.org/10.1016/j.jplph.2014.07.027>
- HAUGWITZ, M.S. et al. (2014). Soil microorganisms respond to five years of climate change manipulations and elevated atmospheric CO₂ in a temperate heath ecosystem. *Plant Soil*, 374, 211–222. <https://doi.org/10.1007/s11104-013-1855-1>
- HUANG, X. et al. (2014). Changes of soil microbial biomass carbon and community composition through mixing nitrogen-fixing species with *Eucalyptus urophylla* in subtropical China. *Soil Biol. Biochem.*, 73, 42–48. <https://doi.org/10.1016/j.soilbio.2014.01.021>
- INEICHEN, K. WIEMKEN, V. and WIEMKEN, A. (1995). Shoots, roots and ectomycorrhiza formation of pine seedlings at elevated atmospheric carbon dioxide. *Plant Cell Environ.*, 18, 703–707.
- IPCC (2013). Summary for Policymakers. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. New York, NY: Cambridge University Press.
- JIN, J. et al. (2013). Elevated CO₂ temporally enhances phosphorus immobilization in the rhizosphere of wheat and chickpea. *Plant Soil*, 368, 315–328. <https://doi.org/10.1007/s11104-012-1516-9>
- JOHNSON, N.C. et al. (2013). Predicting community and ecosystem outcomes of mycorrhizal responses to global change. *Ecology Letters*, 16(Suppl. 1), 140–153. <https://doi.org/10.1111/ele.12085>
- JOHNSON, N.C. et al. (2005). Species of plants and associated arbuscular mycorrhizal fungi mediate mycorrhizal responses to CO₂ enrichment. *Global Change Biology*, 11, 1156–1166.
- JOHNSON, N.C. and GEHRING, C.A. (2007). Mycorrhizas: symbiotic mediators of rhizosphere and ecosystem processes. In: Cardon, Z.G., Whitbeck, J.L. (Eds.). *The Rhizosphere: An Ecological Perspective*. London: Elsevier Academic Press (pp. 31–56).
- KABIR Z. et al. (1997). Seasonal changes of arbuscular mycorrhizal fungi as affected by tillage practices and fertilization: hyphal density and mycorrhizal root colonization. *Plant Soil*. 192(2), 285–293. <https://doi.org/10.1023/A:1004205828485>
- KUMAR, A. et al. (2019). Effects of water deficit stress on agronomic and physiological responses of rice and greenhouse gas emission from rice soil under elevated atmospheric CO₂. *Sci. Total Environ.*, 650, 2032–2050
- KUZYAKOV, Y. et al. (2018). Review and synthesis of the effects of elevated atmospheric CO₂ on soil processes: No changes in pools, but increased fluxes and accelerated cycles. *Soil Biology and Biochemistry*, 128, 66–78. <https://doi.org/10.1016/j.soilbio.2018.10.005>
- LIU, S. et al. (2018). Climatic role of terrestrial ecosystem under elevated CO₂: a bottom-up greenhouse gases budget. *Ecology Letters*, 21(7), 1108–1118. <https://doi.org/10.1111/ele.13078>
- MATAMALA, R. and DRAKE, B.G. (1999). The influence of atmospheric CO₂ enrichment on plant–soil nitrogen interactions in a wetland plant community on the Chesapeake Bay. *Plant Soil*, 210, 93–101.
- MCCARTHY, H.R. et al. (2010). Re-assessment of plant carbon dynamics at the Duke free-air CO₂ enrichment site: Interactions of atmospheric CO₂ with nitrogen and water availability over stand development. *New Phytologist*, 185(2), 514–528. <https://doi.org/10.1111/j.1469-8137.2009.03078.x>
- MORAN, K.K. and JASTROW, J.D. (2010). Elevated carbon dioxide does not offset loss of soil carbon from a corn-soybean agroecosystem. *Environmental Pollution*, 158(4), 1088–1094. <https://doi.org/10.1016/j.envpol.2009.07.005>
- NIE, M. et al. (2013). Positive climate feedbacks of soil microbial communities in a semi-arid grassland. *Ecol. Lett.*, 16(2), 234–241. <https://doi.org/10.1111/ele.12034>
- OLIVEIRA V.F. et al (2010). Elevated CO₂ atmosphere promotes plants growth and inulin production in the cerrado species *Vernonia herbacea*. *Funct Plant Biol*, 37, 223–231.
- PANNEERSELVAM P. et al. (2019). Influence of elevated CO₂ on arbuscular mycorrhizal fungal community elucidated using Illumina MiSeq platform in sub-humid tropical paddy soil. *Applied Soil Ecology*, 145, 103344, 9. <https://doi.org/10.1016/j.apsoil.2019.08.006>
- PENDALL, E. et al. (2013). Warming reduces carbon losses from grassland exposed to elevated atmospheric carbon dioxide. *PLoS One*, 8, e71921. <https://doi.org/10.1371/journal.pone.0071921>

- PHILLIPS, J.M. and HAYMAN D.S. (1970). Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans. British Mycology Society*, 55, 150–160.
- PROCTER, A.C. et al. (2014). Fungal community responses to past and future atmospheric CO₂ differ by soil type. *Appl. Environ. Microbiol.*, 80, 7364–7377. <https://doi.org/10.1128/AEM.02083-14>
- REINSCH, S. et al. (2013). Impact of future climatic conditions on the potential for soil organic matter priming. *Soil Biol. Biochem.*, 65, 133–140. <https://doi.org/10.1016/j.soilbio.2013.05.013>
- RILLIG, M.C. and ALLEN, M.F. (1999). What is the role of arbuscular mycorrhizal fungi in plant-to-ecosystem responses to elevated atmospheric CO₂? *Mycorrhiza*, 9, 1–8.
- ROGERS A, Y. et al. (2006). Increased C availability at elevated carbon dioxide concentration improves N assimilation in a legume. *Plant, Cell and Environment*, 29, 1651–1658.
- SAITOH Y et al. (2004). Yeast generated CO₂ as a convenient source of CO₂ for adult mosquito sampling. *Journal of American Mosquitoes Control Association*, 20, 261–264.
- SAKARIYAWO O.S. et al. (2016). Growth, assimilate partitioning and grain yield response of soybean (*Glycine max* L. Merrill) varieties to carbon dioxide enrichment and arbuscular mycorrhizal fungi in the humid rainforest. *Agro-science*, 15(2), 29–40. <https://doi.org/10.4314/as.v15i2.5>
- SCHORTEMAYER, M. et al. (2000). Effects of elevated atmospheric CO₂ concentration in C and N pools and rhizosphere processes in a Florida scrub oak community. *Glob. Change Biol.*, 6, 383–391.
- SILLEN, W.M.A. and DIELEMAN W.I.J. (2012). Effects of elevated CO₂ and N fertilization on plant and soil carbon pools of managed grasslands: a meta-analysis. *Biogeosciences*, 9, 2247–2258. <https://doi.org/10.5194/bg-9-2247-2012>
- SINGH, S.B. et al. (2017). Impact of secondary forest fallow period on soil microbial biomass carbon and enzyme activity dynamics under shifting cultivation in North Eastern Hill region, India. *Catena*, 156, 10–17. <https://doi.org/10.1016/j.catena.2017.03.017>
- SINGH, L.P., GILL, S.S. and TUTEJA, N. (2011). Unravelling the role of fungal symbionts in plant abiotic stress tolerance. *Plant Signal. Behav.*, 6(2), 175–191. <https://dx.doi.org/10.4161/psb.6.2.14146>
- SMITH, S.E. and READ, D.J. (2008). *Mycorrhizal Symbiosis*. 3rd ed. London: Academic Press (pp. 316e319).
- VANCE E.D. (1987). An extraction method for measuring soil microbial biomass C. *Soil Biol. Biochem.*, 19, 703–707. [https://doi.org/10.1016/0038-0717\(87\)90052-6](https://doi.org/10.1016/0038-0717(87)90052-6)
- WEIGEL, H.J. et al. (2005). Carbon turnover in a crop rotation under free air CO₂ enrichment (FACE). *Pedosphere*, 15, 728–738.
- WIPF, D. et al. (2019). Trading on the arbuscular mycorrhiza market: from arbuscules to common mycorrhizal networks. *New Phytol.*, 223(3), 1127–1142. <https://doi.org/10.1111/nph.15775>
- WU, D.X. et al. (2004). Effects of elevated CO₂ concentration on growth, water use, yield and grain quality of wheat under two soil water levels. *Agriculture, Ecosystems & Environment*, 104(3), 493–507. <https://doi.org/10.1016/j.agee.2004.01.018>
- YANG, L. et al. (2006). Seasonal changes in the effects of free-air CO₂ enrichment (FACE) on dry matter production and distribution of rice (*Oryza sativa* L.). *Field Crops Research*, 98, 12–19. <https://doi.org/10.1016/j.fcr.2005.11.003>
- ZISKA, L. H. (2000). The impact of elevated CO₂ on yield loss from a C₃ and C₄ weed in field-grown soybean. *Global Change Biology*, 6, 899–905.